



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Nystagmus

Tarnutzer, Alexander A ; Straumann, Dominik

DOI: <https://doi.org/10.1097/WCO.0000000000000517>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-141968>

Journal Article

Published Version

Originally published at:

Tarnutzer, Alexander A; Straumann, Dominik (2018). Nystagmus. Current Opinion in Neurology, 31(1):74-80.

DOI: <https://doi.org/10.1097/WCO.0000000000000517>



Nystagmus

Alexander A. Tarnutzer^{a,b} and Dominik Straumann^{a,b}

Purpose of review

The clinical and laboratory assessment of nystagmus in patients with neurologic disorders can provide crucial elements for a state-of-the-art differential diagnosis. An increasing number of publications in the fields of neuro-otology and neuro-ophthalmology have nystagmus in the center of interest, which makes frequent updates on the diagnostic and therapeutic relevance of these contributions indispensable. This review covers important clinical studies and studies in basic research relevant for the neurologist published from January 2016 to August 2017.

Recent findings

Current themes include vestibular nystagmus, positional nystagmus, optokinetic nystagmus and after-nystagmus, vibration-induced nystagmus, head-shaking nystagmus, postrotatory nystagmus, caloric nystagmus, nystagmus in cerebellar disorders, differential diagnosis of nystagmus and treatment approaches (whereas infantile nystagmus syndrome is not addressed in this review). These studies address mechanisms/pathomechanisms, differential diagnoses and treatment of different forms of nystagmus.

Summary

In clinical practice, a structured description of nystagmus including its three-dimensional beating direction, trigger factors and duration is of major importance. The differential diagnosis of downbeat nystagmus is broad and includes acute intoxications, neurodegenerative disorders and cerebrovascular causes amongst others. In patients with positional nystagmus, the distinction between frequent benign peripheral and rare but dangerous central causes is imperative.

Keywords

cerebellum, downbeat nystagmus, eye movements, positional nystagmus, vestibulo-ocular reflex

INTRODUCTION

Nystagmus refers to repetitive, to-and-fro movements of the eyes that are initiated by slow phases [1]. Different types of nystagmus can be distinguished based on the presence of fast and slow phases (jerk nystagmus) or alternating slow phases (pendular nystagmus). In the assessment of nystagmus, a structured approach is essential, as the differential diagnosis of nystagmus is very broad. This includes both benign peripheral-vestibular disorders and dangerous potentially life-threatening central causes. In this review, we summarize recent advances in the assessment of nystagmus in both healthy human individuals and patients. Separate sections will address different forms of nystagmus, including vertical nystagmus, positional nystagmus, head-shaking and vibration-induced nystagmus and vestibular nystagmus. Note that this review does not address publications on infantile nystagmus syndrome (congenital nystagmus).

VESTIBULAR NYSTAGMUS

Of major importance is the distinction between peripheral-vestibular and central nystagmus. The presence of anticompensatory eye movements (AQEM) during video head impulse testing was proposed to be predictive for a peripheral-vestibular origin of spontaneous nystagmus compared with central causes [2]. Recently, nystagmus was assessed during Menière attacks. Although the beating direction was towards the affected side in two patients with unilateral Menière's disease, suggesting an irritative nystagmus in the acute phase [3], ictal

^aDepartment of Neurology, University Hospital Zurich and ^bUniversity of Zurich, Zurich, Switzerland

Correspondence to Dominik Straumann, MD, Department of Neurology, University Hospital Zurich, Frauenklinikstr. 26, CH-8091 Zurich, Switzerland. Tel: +41 44 255 11 11; fax: +41 44 255 43 80; e-mail: dominik.straumann@usz.ch

Curr Opin Neurol 2017, 30:000–000

DOI:10.1097/WCO.0000000000000517

KEY POINTS

- A structured description of nystagmus including its three-dimensional beating direction, trigger factors and duration is recommended.
- In patients with positional nystagmus, the distinction between frequent benign peripheral and rare but dangerous central causes is imperative.
- The differential diagnosis of downbeat nystagmus is broad and includes acute intoxications, neurodegenerative disorders and cerebrovascular causes amongst others.

downbeat nystagmus (DBN) was reported in a patient with bilateral Menière's disease, possibly because of posterior canal inhibition [4]. Whereas pure vertical nystagmus is typically of central origin, bilateral superior canal plugging in two patients with bilateral superior canal dehiscence resulted in vibration-induced upbeat nystagmus, reflecting an unusual peripheral cause for upbeat nystagmus [5].

POSITIONAL NYSTAGMUS

Positional vertigo and nystagmus may present a diagnostic challenge. Although most frequently observed in benign paroxysmal positional vertigo (BPPV), central disorders may result in positional nystagmus as well. Rates of asymptomatic positional nystagmus in healthy controls reach almost 90% [6]. A recent systematic review on clinical and radiological findings in lesion-induced central positional nystagmus concluded that better phenotyping of central positional nystagmus is needed to allow a reliable discrimination from peripheral causes [7[¶]].

The bow and lean test can be useful in identifying the type (geotropic versus apogeotropic variant) and side (right versus left) of lateral canal BPPV. In a recent study, evaluating nystagmus intensity and direction with the head bent forward ('bow') and backward ('lean'), correct identification was achieved in 79% of patients [8[¶]]. Noteworthy, positional nystagmus may show a change in beating direction over time. Short-term adaptation of the vestibulo-ocular reflex has been proposed to explain the formation of direction-reversing nystagmus, which was found more frequently in lateral canal BPPV compared with posterior canal BPPV [9]. Paroxysmal positional nystagmus may also be associated with a fistula between the lateral canal deformity and an enlarged vestibular aqueduct [10].

In supine position, lateral canal BPPV may mimic spontaneous nystagmus. Such "pseudo-

spontaneous' nystagmus was more frequent in patients with lateral canal cupulolithiasis than canalolithiasis (93 vs. 41%) and is likely mediated by the natural inclination of the lateral canal and the medial-to-lateral orientation of the cupular axis in cupulolithiasis and by direction-reversing nystagmus in canalolithiasis [11].

Persistent geotropic direction-changing positional nystagmus (DCPN) with neutral position and cupulolithiasis of the lateral canal was recently characterized by a short clinical course and frequent recurrence [12]. Noteworthy, those patients with long-duration geotropic DCPN did not benefit from canalith-repositioning procedures, suggesting that the underlying disorder is not linked to free-floating debris but rather to deflection of the cupula [13].

New insights into mechanisms of the light cupula syndrome were described in two case reports: whereas heavy endolymph was proposed as a mechanism for persistent positional vertigo and geotropic positional nystagmus in a patient with meningitis and high cerebrospinal fluid (CSF) protein [14], a light cupula of a posterior canal was suggested in a case with persistent torsional DBN [15]. Novel therapeutic strategies in geotropic DCPN include treatment with transcutaneous vagus nerve stimulation [16], that led the authors to propose a top-down mechanism that normalized lateral canal hyperexcitability. In patients with vestibular migraine and positional vertigo, prophylactic treatment with cinnarizine or topiramate resulted in cessation of symptoms in 92% of patients [17].

OPTOKINETIC NYSTAGMUS AND AFTER-NYSTAGMUS

Optokinetic nystagmus (OKN) is elicited by full-field visual movement, whereby both the smooth pursuit system (via the foveal retina) and the optokinetic system (via the extra-foveal retina) are stimulated. The contribution of the optokinetic system to the overall ocular motor response during full-field visual movement is reflected in the optokinetic after-nystagmus (OKAN), which can be observed whenever OKN is terminated by a sudden switch to total darkness. The gain of the optokinetic nystagmus is defined by dividing the velocity of the nystagmus slow phase by the velocity of the visual stimulus. The generation of OKN fast phases is a biological time process that can be influenced by a concurrent cognitive task [18[¶]]. Measuring the optokinetic response may be used as an objective measure of contrast sensitivity function [19] or visual acuity [20].

OKN can be suppressed by looking at a space-fixed fixation light. If, during that task, patients have to detect the appearance of additional targets

without looking at them, the OKN is less suppressed and the effect of this divided attention is more pronounced if the additional target is moving [21]. Whenever oppositely directed optokinetic stimuli are presented simultaneously in different areas of the visual field, attention to one of the fields increases the gain of the OKN in the direction of the visual stimulus in this field [22].

The time constant of the OKAN reflects the effectiveness of the velocity-storage mechanism, which, in turn, plays an important role in motion sickness [23]. Accordingly, a strong positive correlation between the OKAN time constant and the susceptibility to visually induced motion sickness was found [24].

Whenever a horizontal optokinetic stimulus is projected for 2 s on the two eyes, but with an opposite movement direction, the OKN and the perceived movement point in the same direction in both healthy individuals and patients with Parkinson's disease; thus, OKN in this paradigm can be used to indicate conscious motion perception [25].

VIBRATION-INDUCED NYSTAGMUS

Vibration-induced nystagmus is elicited by pressing a vibrating device perpendicularly to the skin over the mastoid process, typically during 20 s. In patients with a unilateral vestibular hypofunction, this bone-conducted vibration induces predominantly horizontal nystagmus in the direction of the unaffected ear. In some patients with Menière's disease or vestibular schwannoma, however, vibration-induced nystagmus may also be directed towards the affected ear [26].

Vibration-induced nystagmus is time-locked with the stimulus, should be observed under Frenzel or infrared-video goggles, and is most robust at a stimulus frequency of 100 Hz [27]. Previously, the most common used frequencies were between 30 and 60 Hz [28]. On the basis of measurements with piezo-electric and force sensors, vibration is more effective on the mastoid than on the posterior cervical muscles and more effective if the apparatus is held by the dominant hand of the examiner [29]. In patients with vestibular schwannoma, nystagmus induced by vibration of the ipsilateral sterno-cleido-mastoid muscle correlated with tumor size [30].

HEAD-SHAKING NYSTAGMUS

Head-shaking nystagmus, or, more precisely, nystagmus after horizontal head shaking, can be observed in patients with peripheral or central asymmetries of vestibular signals between the

two sides. Typically, head-shaking nystagmus is horizontal. In some patients with central lesion, head-shaking nystagmus may be vertical. This nystagmus is called perverted head-shaking nystagmus. Patients with perverted head-shaking nystagmus frequently show impaired tilt suppression of postrotatory nystagmus [31]. Also, there is an association between perverted head-shaking nystagmus and essential tremor [32].

Perverted head-shaking nystagmus is usually downbeat, rarely upbeat, and a correlation with specific MR-lesions was not found so far [33]. Considering the three-dimensional angular velocity axis and the slow-phase velocity time constant, perverted downbeat head-shaking nystagmus is probably the result of increased activity of the central anterior canal pathways [34].

POSTROTATORY NYSTAGMUS

Postrotatory nystagmus is a vestibular nystagmus elicited by rapid deceleration of a turntable from constant-velocity to zero-velocity rotation about an earth-vertical axis. Typically, patients are in an upright position, and the axis of deceleration passes through the center of the inter-aural line. The slow-phase velocity of horizontal postrotatory nystagmus in the dark decays with a time constant (usually between 5 and 10 s) determined by the velocity-storage mechanisms of the brainstem and cerebellum. Tilting the head or whole body immediately or a few moments after the end of deceleration suppresses postrotatory nystagmus and therefore, decreases the time constant of the slow-phase velocity decay. In addition, the angular slow-phase velocity axis of the eyes shifts in the direction of the gravity vector. In some patients, this tilt suppression of postrotatory nystagmus causes vertigo, disorientation or motion sickness because of the conflict of signals from the semicircular canals and the otoliths. Patients with impaired tilt suppression of postrotatory nystagmus frequently show perverted head-shaking nystagmus [31].

Tilt suppression of postrotatory nystagmus of patients with vestibular migraine differs from healthy individuals and patients with migraine headaches: the angular slow-phase velocity axis of the eyes shows a larger shift in the direction of the gravity vector, but this effect is not correlated with the shortening of the slow-phase velocity time constant [35]. Whether the resulting 'residual sensory conflict' explains the motion sickness susceptibility of the patients with vestibular migraine needs to be further explored [36].

Whenever postrotatory nystagmus is not modified by a tilt, but by centrifugation without an

angular movement (so-called double centrifugation), the time constant of the slow-phase velocity decay is not affected [37]. Thus, otolith signals alone – in this case oscillating inter-aural and naso-occipital translatory signals – do not interfere with the velocity-storage mechanism.

To simulate the effect of horizontal spontaneous nystagmus on the high-acceleration vestibulo-ocular reflex, healthy individuals were assessed with the video head impulse test. It was found that the slow-phase velocity of postrotatory nystagmus and the velocity of the head-impulse evoked eye movements simply added; thus, to compute the correct gain of the vestibulo-ocular reflex during head impulses, one should subtract the velocity of the baseline eye drift [38].

CALORIC NYSTAGMUS

Irrigation of the external auditory canal with warm or cold water elicits a mainly horizontal nystagmus, so-called caloric nystagmus. As the cold or warm temperature from the external auditory canal reaches the lateral semicircular canal through the mastoid bone, the individual anatomy of this bone could determine the intensity of the evoked caloric nystagmus. In a preliminary study, it was found that the amount of airspace in the mastoid bone is inversely related to the effectiveness of warm caloric irrigation [39]. Mental tasking, on the other side, does not influence the parameters of caloric nystagmus [40].

Strong cold-water irrigation with 24 °C reduces the gain of the vestibulo-ocular reflex during

ipsilateral head impulses, whereas weaker cold-water or warm-water irrigation has no effect on the result of the head impulse test [41]. The strong cold-water irrigation may shift the set point of the nonlinear relation between head acceleration and the vestibular firing rate toward a less acceleration-sensitive zone, thus, demonstrating Ewald's second law in healthy individuals.

NYSTAGMUS IN CEREBELLAR DISORDERS

To better understand the impact of vertical nystagmus in patients, knowing its prevalence in the general population is important. A recently developed nystagmus-specific quality-of-life questionnaire can help to determine the impact of nystagmus on daily living [42]. In a recent study, spontaneous upbeat nystagmus in supine position was noted in almost half of all healthy participants and was linked to individual structural differences in cerebellar tonsil volume [43]. Positional DBN and perverted head-shaking nystagmus seem to be associated with essential tremor, as these findings were more prevalent than in controls (21.2 vs. 3.9%) [32]. Acute-onset nystagmus may be seen in cases with drug intoxication (as, e.g. acute DBN in oxcarbazepine intoxication, see Fig. 1 [44]) or pesticides (avermectins [45]). However, also therapeutic doses of commonly prescribed drugs may trigger DBN and dizziness, as reported for pregabalin [46] and lithium [47], being reversible after cessation in both cases. Blocking of the excitatory responses of vestibular nuclei neurons may explain DBN because of a




FIGURE 1. Electroencephalography (EEG) recording in longitudinal bipolar montage (10/20 system). Panel A: at the time of oxcarbazepine intoxication (day 3), steep surface-negative excursions (arrows) are followed by more slow return to baseline. Considering eyeball polarity (positive to negative from ventral to caudal), this indicates slow upward eye-drift followed by fast downward saccades, consistent with DBN. Note that eyelid movements will contribute to these EEG changes as well. Panel B: 48 h later DBN had resolved, consistent with the return of oxcarbazepine blood levels to the therapeutic range. This figure was originally published in Ref. [44].




FIGURE 2. Blood alcohol content (BAC) on eye-drift velocity as a function of eye gaze angle. Effect of 0.6‰ BAC on eye-drift velocity as a function of eye gaze angle. Each line represents the mean drift velocity of all individuals in the different conditions, whereas the shaded area represents the mean (SD). The light grey dashed line is a scaled version of the data recorded before alcohol intake (dark grey dashed line), perfectly overlapping with the data recorded after alcohol intake (black solid line), confirming the pure scaling effect of 0.6‰ BAC. Such an effect is further confirmed by the scaling parameter of the tangent model (k_2), which was estimated on the plotted curves (the estimated parameters are reported in the figure). This figure was originally published in Ref. [49[¶]].

selective histamine H2 receptor antagonist (ranitidine) [48].

As a novel human model to study effects of cerebellar loss-of-function on gaze-holding properties, controlled alcohol intake was proposed [49[¶]]. At a blood-alcohol content of 0.06%, horizontal eye-drift velocity approximately doubled at all eccentricities studied (Fig. 2), matching the pattern of gaze-holding deficits previously noted in late-onset neurodegenerative cerebellar disease.

During the diagnostic procedure for new-onset nystagmus, caution is advised for patients with apogeotropic positional nystagmus during supine-roll testing whenever repeated repositioning maneuvers remain without success. In these patients, midline cerebellar mass lesions may mimic BPPV [50].

NYSTAGMUS IN THE DIFFERENTIAL DIAGNOSIS

The localizing value of ocular motor findings was emphasized in a patient with conjugate ipsilesional

eye deviation and spontaneous horizontal nystagmus beating contralesionally in lateral medullary infarction [51]. Sub-acute onset of upbeat nystagmus and gaze-evoked nystagmus in combination with neurocognitive deficits and neuropathy after bariatric surgery should raise a high suspicion of Wernicke encephalopathy and prompt immediate treatment with parenteral thiamine [52].

Acquired periodic alternating nystagmus (PAN) was linked to autoimmune disorders and long-term lithium intake, promoting the application of intravenous immunoglobulin in the first case [53] and prescription of baclofen in the latter case [54]. In a patient with systemic lupus erythematosus and cerebellar atrophy, amantadine reduced PAN, dizziness and oscillopsia [55]. Lacking central (cerebellar) adaptive recalibration because of profound visual loss has been proposed to explain windmill nystagmus, a rare type of nystagmus that is characterized by a clock-like rotation of the nystagmus beating direction [56].

Ocular motor cerebellar signs and ataxia of gait/stance may precede classical features in sporadic Creutzfeldt–Jakob disease [57] and may reflect prominent features in immune-mediated anti-DPPX encephalitis [58]. Hereditary cerebellar neurodegeneration may initially present as (positional) DBN, as described for spinocerebellar ataxia type 6 [59] and X-linked adrenoleukodystrophy [60]. Hypometric horizontal saccades, disconjugate gaze-evoked nystagmus and esotropia may represent prominent features in central nervous system Whipple's disease [61].

In children, vertical nystagmus may be linked to retinal dystrophies [62]. Different nystagmus patterns may be seen in newborn and infants with Joubert syndrome [63], mutations in the calcium/calmodulin-dependent serine protein kinase (CASK) gene leading to FG-syndrome 4 [64], visual deprivation from child neglect [65], mass lesions above the midbrain [66] and galactokinase deficiency [67]. The differential diagnosis of isolated nystagmus in pediatric patients remains challenging. In a case series of 148 children with isolated nystagmus, only 23 (15.5%) had abnormalities on brain MRI, with abnormal signal lesions, Chiari type 1 malformations and optic glioma most frequently reported [68].

TREATMENT APPROACHES TO NYSTAGMUS AND OSCILLOPSIA

A broad range of drugs has been investigated regarding their effect on different forms of nystagmus [69[¶]]. For channelopathies as familial hemiplegic migraine, acetazolamide is effective for treating

disequilibrium and positioning DBN [70]. In autoimmune-mediated glutamic acid decarboxylase (GAD)-associated cerebellar ataxia, intravenous immunoglobulin successfully treated paroxysmal positioning upbeat nystagmus [71].

Novel noninvasive therapeutic approaches to reduce DBN include real-time computer-based visual feedback, reducing slow-phase vertical eye movements and improving visual acuity in 5 of 10 patients [72] and resting in darkness for 2 h (compared with remaining in bright light), resulting in a relative reduction of vertical slow-phase eye velocity by about one-third [73]. Advanced therapeutic interventions in acquired nystagmus include the attachment of a T-plate to the lateral orbital rim and the inferior rectus muscle tendon [74], the implantation of an oculomotor prosthesis [75] and retrobulbar botulinum-toxin A injections [76], resulting in reduced nystagmus amplitude and oscillopsia in all cases. Lateral canal occlusion was confirmed as a valuable treatment option for head-shaking nystagmus and vertigo in a patient with head-jolting nystagmus syndrome [77].

CONCLUSION

Nystagmus is a frequent sign in patients with neurological disorders. Physiological nystagmus includes rotatory and postrotatory nystagmus, OKN and after-nystagmus, as well as symmetric caloric nystagmus. Pathological nystagmus includes spontaneous nystagmus, positional nystagmus, vibration-induced nystagmus, head-shaking nystagmus and asymmetric caloric nystagmus. The differential diagnosis and therapy of pathological nystagmus relies on an increasing body of information detailing the mechanisms or pathomechanisms of different forms of nystagmus.

Acknowledgements

None.

Financial support and sponsorship

None.

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Leigh RJ, Zee DS. The neurology of eye movements. USA: Oxford University Press; 2015.
 2. Luis L, Lehnen N, Muñoz E, *et al.* Anticompensatory quick eye movements after head impulses: a peripheral vestibular sign in spontaneous nystagmus. *J Vestib Res* 2016; 25:267–271.
 3. Hirai C, Yamamoto Y, Takeda T, *et al.* Nystagmus at the onset of vertiginous attack in Menière's disease. *Otol Neurotol* 2017; 38:110–113.
 4. Lee S-U, Kim H-J, Lee E-S, *et al.* Ictal downbeat nystagmus in bilateral Menière's disease. *J Neurol* 2017. [Epub ahead of print]
 5. Sharon JD, Carey JP, Schubert MC. Upbeat nystagmus after bilateral superior canal plugging: a peripheral cause of vertical nystagmus. *Laryngoscope* 2017; 127:1698–1700.
 6. Martens C, Goplen FK, Nordfalk KF, *et al.* Prevalence and characteristics of positional nystagmus in normal subjects. *Otolaryngol Head Neck Surg* 2016; 154:861–867.
 7. Macdonald NK, Kaski D, Saman Y, *et al.* Central positional nystagmus: a systematic literature review. *Front Neurol* 2017; 8:141.
- This systematic review summarizes with great detail current knowledge in central positional nystagmus and identifies areas of uncertainty. It emphasizes the need for further research on central positional nystagmus.
8. Marcelli V. Nystagmus intensity and direction in bow and lean test: an aid to diagnosis of lateral semicircular canal benign paroxysmal positional vertigo. *Acta Otorhinolaryngol Ital* 2016; 36:520–526.
- This study provides a three-step approach to lateral canal BPPV which will help the clinician to choose the correct repositioning maneuver.
9. Jeong K-H, Shin JE, Shin DH, Kim C-H. Direction-reversing nystagmus in horizontal and posterior semicircular canal canalolithiasis. *Otol Neurotol* 2016; 37:767–771.
 10. Suzuki M, Ota Y, Tanaka T, Ota Y. Lateral semicircular canal-enlarged vestibular aqueduct fistula associated with paroxysmal positional nystagmus. *Otol Neurotol* 2016; 37:e192–e193.
 11. Im DH, Yang YS, Choi H, *et al.* Pseudo-spontaneous nystagmus in horizontal semicircular canal canalolithiasis. *Medicine (Baltimore)* 2017; 96:e7849.
 12. Seo T, Shiraiishi K, Kobayashi T, *et al.* Clinical course of persistent geotropic direction-changing positional nystagmus with neutral position - Light cupula. *Acta Otolaryngol* 2016; 136:34–37.
 13. Ban JH, Kim M-B, Hong SM. Immediate and short-term therapeutic results between direction-changing positional nystagmus with short- and long-duration groups. *Ear Hear* 2016; 37:243–246.
 14. Choi J-Y, Lee E-S, Kim H-J, Kim JS. Persistent geotropic positional nystagmus after meningitis: evidence for light cupula. *J Neurol Sci* 2017; 379:279–280.
 15. Ichimura A, Otsuka K. Persistent down-beating torsional positional nystagmus: posterior semicircular canal light cupula? *Case Rep Otolaryngol* 2016; 2016:1249325.
 16. Cha WW, Song K, Lee HY. Persistent geotropic direction-changing positional nystagmus treated with transcutaneous vagus nerve stimulation. *Brain Stimul* 2016; 9:469–470.
 17. El-Badry MM, Samy H, Kabel AM, *et al.* Clinical criteria of positional vertical nystagmus in vestibular migraine. *Acta Otolaryngol* 2017; 137:720–722.
 18. Balaban CD, Furman JM. Beat-to-beat control of human optokinetic nystagmus slow phase durations. *J Neurophysiol* 2017; 117:204–214.
- This study demonstrates that the timing of optokinetic fast phases can be influenced by cognitive processes.
19. Dakin SC, Turnbull PRK. Similar contrast sensitivity functions measured using psychophysics and optokinetic nystagmus. *Sci Rep* 2016; 6:34514.
 20. Bögli SY, Afthinos M, Bertolini G, *et al.* Unravelling stimulus direction dependency of visual acuity in larval zebrafish by consistent eye displacements upon optokinetic stimulation. *Invest Ophthalmol Vis Sci* 2016; 57:1721–1727.
 21. Williams IM, Schofield P, Khade N, Abel LA. Divided visual attention: a comparison of patients with multiple sclerosis and controls, assessed with an optokinetic nystagmus suppression task. *J Clin Neurosci* 2016; 34:187–192.
 22. Kanari K, Sakamoto K, Kaneko H. Effect of visual attention on the properties of optokinetic nystagmus. *PLoS One* 2017; 12:e0175453.
 23. Dai M, Kunin M, Raphan T, Cohen B. The relation of motion sickness to the spatial-temporal properties of velocity storage. *Exp Brain Res* 2003; 151:173–189.
 24. Guo CCT, Chen DJZ, Wei IY, *et al.* Correlations between individual susceptibility to visually induced motion sickness and decaying time constant of after-nystagmus. *Appl Ergon* 2017; 63:1–8.
 25. Fujiwara M, Ding C, Kaunitz L, *et al.* Optokinetic nystagmus reflects perceptual directions in the onset binocular rivalry in Parkinson's disease. *PLoS One* 2017; 12:e0173707.
 26. Vargas Gamarra MF, Krstulovic C, Pérez Guillén V, Pérez Garrigues H. Ipsilesional nystagmus induced by vibration in subjects with Ménière's disease or vestibular schwannoma. *Otol Neurotol* 2017; 38:e168–e172.
 27. Dumas G, Curthoys IS, Lion A, *et al.* The skull vibration-induced nystagmus test of vestibular function-a review. *Front Neurol* 2017; 8:41.
 28. Hamann K-F. Vibration-Induced Nystagmus: A Biomarker for Vestibular Deficits - A Synopsis. *ORL J Otorhinolaryngol Relat Spec* 2017; 79:112–120.
 29. Dumas G, Lion A, Perrin P, *et al.* Topographic analysis of the skull vibration-induced nystagmus test with piezoelectric accelerometers and force sensors. *Neuroreport* 2016; 27:318–322.

30. Lee JM, Kim MJ, Kim JW, *et al.* Vibration-induced nystagmus in patients with vestibular schwannoma: characteristics and clinical implications. *Clin Neurophysiol* 2017; 128:1372–1379.
31. Lee S-U, Choi J-Y, Kim H-J, *et al.* Impaired tilt suppression of post-rotatory nystagmus and cross-coupled head-shaking nystagmus in cerebellar lesions: image mapping study. *Cerebellum* 2017; 16:95–102.
32. Kim YE, Kim JS, Yang H-J, *et al.* Perverted head-shaking and positional downbeat nystagmus in essential tremor. *Cerebellum* 2016; 15:152–158.
33. Han W-G, Yoon H-C, Kim T-M, *et al.* Clinical correlation between perverted nystagmus and brain MRI abnormal findings. *J Audiol Otol* 2016; 20:85–89.
34. Choi J-Y, Jung I, Jung J-M, *et al.* Characteristics and mechanism of perverted head-shaking nystagmus in central lesions: video-oculography analysis. *Clin Neurophysiol* 2016; 127:2973–2978.

This study provides experimental evidence that perverted head-shaking nystagmus may be due to increased activity of the central anterior canal pathways.

35. Wang J, Lewis RF. Contribution of intravestibular sensory conflict to motion sickness and dizziness in migraine disorders. *J Neurophysiol* 2016; 116:1586–1591.
36. Jeong S-H, Kim JS. Direction and magnitude of head tilt alter postrotatory nystagmus. *J Neurophysiol* 2016; 116:2000–2010.
37. Shaikh AG, Solomon D. Effects of sustained otolith-only stimulation on post-rotational nystagmus. *Cerebellum* 2017; 16:683–690.
38. Mantokoudis G, Tehrani ASS, Xie L, *et al.* The video head impulse test during postrotatory nystagmus: physiology and clinical implications. *Exp Brain Res* 2016; 234:277–286.
39. Patki AU, Ronen O, Kaylie DM, *et al.* Anatomic variations in temporal bones affect the intensity of nystagmus during warm caloric irrigation. *Otol Neurotol* 2016; 37:1111–1116.

This preliminary study explains the intensity of caloric nystagmus by the amount of airspace in the mastoid bone.

40. Easterday MK, Plyler PN, Doettl SM. Mental tasking and caloric-induced vestibular nystagmus utilizing videonystagmography. *Am J Audiol* 2016; 25:177–183.
41. Tamás LT, Weber KP, Bockisch CJ, *et al.* Cold thermal irrigation decreases the ipsilateral gain of the vestibulo-ocular reflex. *Ear Hear* 2017; 38:e193–e199.
42. McLean RJ, Maconachie GDE, Gottlob I, Maltby J. The development of a nystagmus-specific quality-of-life questionnaire. *Ophthalmology* 2016; 123:2023–2027.
43. Ruehl RM, Stephan T, Dieterich M, Eulenburg PZ. Voxel-based morphometry delineates the role of the cerebellar tonsil in physiological upbeat nystagmus. *J Neurol* 2017. [Epub ahead of print]

This study provides further evidence for the cerebellar tonsil's role in vertical gaze holding in healthy human individuals.

44. Tarnutzer AA, Imbach LL. The intoxicated EEG. *Seizure* 2016; 45:100–101.
45. Wenjie W, Houqing L, Ping Z, Gengyun S. Nystagmus following acute avermectins poisoning. *Toxicol Ind Health* 2016; 32:751–752.
46. Ekinici AS, Ciftci S, Cavus B, Aydogdu I. Could pregabalin cause oculomotor symptoms in lower dose? A case with down beat nystagmus as a side effect. *Acta Neurol Belg* 2016. [Epub ahead of print]
47. Jørgensen JS, Landschoff Lassen L, Wegener M. Lithium-induced downbeat nystagmus and horizontal gaze palsy. *Open Ophthalmol J* 2016; 10:126–128.
48. Butragueño Laiseca L, Toledo Del Castillo B, Rodríguez Jimenez C, *et al.* Downbeat nystagmus due to ranitidine in a pediatric patient. *Eur J Paediatr Neurol* 2017; 21:682–684.
49. Romano F, Tarnutzer AA, Straumann D, *et al.* Gaze-evoked nystagmus induced by alcohol intoxication. *J Physiol* 2017; 595:2161–2173.

This study proposes a human model for studying transient cerebellar loss of function noninvasively. Although the focus was on gaze holding in the data published, a much broader range of cerebellar impairments may be addressed.

50. Cho B-H, Kim SH, Kim S-S, *et al.* Central positional nystagmus associated with cerebellar tumors: Clinical and topographical analysis. *J Neurol Sci* 2017; 373:147–151.
51. Kobayashi Z, Numasawa Y, Tomimitsu H, Shintani S. Conjugate eye deviation plus spontaneous nystagmus as a diagnostic sign of lateral medullary infarction. *J Neurol Sci* 2016; 367:222–223.
52. Vogrig A, Zannoni T, Moretto G. Nystagmus and lower extremity hyperalgesia after colectomy. *JAMA* 2016; 316:1488–1489.
53. Argente-Escrig H, Bataller L, Krstulovic Roa C, *et al.* Atypical periodic alternating nystagmus responding to high-dose intravenous immunoglobulins: a case report. *J Neuroinflammation* 2017; 14:71.
54. Rust H, Lutz N, Honegger F, *et al.* Periodic alternating nystagmus in a patient on long-term lithium medication. *J Neurol Sci* 2016; 369:252–253.
55. Lee S-H, Lee S-Y, Choi S-M, *et al.* Resolution of periodic alternating nystagmus with amantadine. *J Neurol Sci* 2016; 364:65–67.
56. Choi K-D, Shin HK, Kim JS, *et al.* Variants of windmill nystagmus. *J Neurol* 2016; 263:1375–1381.
57. Choi Y-J, Kang K-W, Lee S-Y, *et al.* Creutzfeldt-Jakob disease presenting with dizziness and gaze-evoked nystagmus: a case report. *Medicine (Baltimore)* 2016; 95:e2766.
58. Doherty L, Gold D, Solnes L, *et al.* Anti-DPPX encephalitis: prominent nystagmus reflected by extraocular muscle FDG-PET avidity. *Neurol Neuroimmunol Neuroinflamm* 2017; 4:e361.
59. Ueno T, Nishizawa H, Suzuki C, *et al.* Downbeat nystagmus as an initial clinical sign in spinocerebellar ataxia type 6. *Neurol Sci* 2017; 38:1543–1545.
60. Kim SH, Kim S-S, Ha H, Lee S-H. X-linked adrenoleukodystrophy presenting with positional downbeat nystagmus. *Neurology* 2016; 86:2214–2215.
61. Shaikh AG, Ghasia FF. Novel eye movement disorders in Whipple's disease-staircase horizontal saccades, gaze-evoked nystagmus, and esotropia. *Front Neurol* 2017; 8:321.
62. Matalia J, Rajput VK, Chillal GJ, Shetty BK. Upbeat nystagmus in a 3.5-year-old boy. *J AAPOS* 2016; 20:88–90.
63. Salva I, Albuquerque C, Moreira A, Dâmaso C. Nystagmus in a newborn: a manifestation of Joubert syndrome in the neonatal period. *BMJ Case Rep* 2016; 2016; pii: bcr2015213127.
64. Dunn P, Prigatano GP, Szelinger S, *et al.* A de novo splice site mutation in CASK causes FG syndrome-4 and congenital nystagmus. *Am J Med Genet A* 2017; 173:611–617.
65. Ho CH, Zargarpour J, Evans M. Raised in darkness: A 7-month-old with nystagmus from severe visual deprivation. *Pediatr Emerg Care* 2017; 33:263–264.
66. İşıkay S. Pendular nystagmus associated with cerebral pyramidal astrocytoma. *Pediatr Neurol* 2016; 61:117–118.
67. Bzduch V, Tomcikova D, Gerinac A, Behulova D. Cataract and early nystagmus due to galactokinase deficiency. *J Inherit Metab Dis* 2017. [Epub ahead of print]
68. Batmanabane V, Héon E, Dai T, *et al.* The role of MR imaging in investigating isolated pediatric nystagmus. *Pediatr Radiol* 2016; 46:1721–1727.
69. Feil K, Bremova T, Muth C, *et al.* Update on the pharmacotherapy of cerebellar ataxia and nystagmus. *Cerebellum* 2016; 15:38–42.

In this review, state-of-the art drug treatment for cerebellar ataxia and nystagmus is summarized.

70. Suzuki M, Fujiwara K, Tsubuku T, *et al.* Time course of downbeat positioning nystagmus in familial hemiplegic migraine type 1 treated with acetazolamide. *J Neurol Sci* 2016; 368:206–208.
71. Martins AI, Carvalho JN, Amorim AM, *et al.* Disabling central paroxysmal positioning upbeat nystagmus and vertigo associated with the presence of anti-glutamic acid decarboxylase antibodies. *J Neuroophthalmol* 2017. [Epub ahead of print]
72. Teufel J, Bardins S, Spiegel R, *et al.* Real-time computer-based visual feedback improves visual acuity in downbeat nystagmus - a pilot study. *J Neuroeng Rehabil* 2016; 13:1.

This pilot-study provides a promising therapeutic approach to DBN.

73. Spiegel R, Claaßen J, Teufel J, *et al.* Resting in darkness improves downbeat nystagmus: evidence from an observational study. *Ann N Y Acad Sci* 2016; 1375:66–73.
74. Tse BC, McKeown CA, Tse DT. Titanium T-plate as a platform for globe stabilization in acquired nystagmus and oscillopsia without a null zone. *Ophthal Plast Reconstr Surg* 2017; 33:e61–e63.
75. Nachev P, Rose GE, Verity DH, *et al.* Magnetic oculomotor prosthetics for acquired nystagmus. *Ophthalmology* 2017; 124:1556–1564.
76. Chen Y-R, Fredrick D, Steinberg GK, Liao YJ. Treatment of nystagmus in brainstem cavernous malformation with botulinum toxin. *Cureus* 2016; 8:e553.
77. Welgampola MS, Croxson GR, McGarvie LA, Halmagyi GM. Head-shaking nystagmus and vertigo cured by lateral semicircular canal occlusion. *J Neurol* 2016; 263:588–590.